ANL/NDM-100

THE ENERGY DEPENDENCE OF THE OPTICAL-MODEL POTENTIAL FOR FAST-NEUTRON SCATTERING FROM BISMUTH

by

A. B. Smith, P. T. Guenther and R. D. Lawson

May 1987

Applied Physics Division Argonne National Laboratory 9700 South Cass Avenue Argonne, Illinois 60439 USA

ABSTRACT

Neutron differential-elastic-scattering cross sections of bismuth were measured at \approx 0.5 MeV intervals from \approx 4.5 to 10.0 MeV. At each incident energy ≥ 40 differential values were obtained, distributed been \approx 180 and 160 . The measured data were combined lower-energy results previously reported from laboratory, and others available in the literature, to provide a detailed data base extending from ≈ 1.5 to 10.0 MeV. This data interpreted in terms the conventional of optical-statistical model and also a model inclusive of the surface-peaked real potential predicted by the dispersion relation. Particular attention was given to the energy dependence of the volume-integral-per-nucleon of the real potential, J., to see if there was evidence of the Fermi Surface In the range 3.0 to 10.0 MeV the present data indicate that dJ,/dE is essentially constant, with a relatively large

negative value of -6.0 to -9.0 fm 3 , depending on the model used in the analysis. Below 3.0 MeV, there is some evidence for a decrease in the magnitude of $\mathrm{dJ_V/dE}$. However, the effect is very small and it is only when this trend is combined with considerations of the J values needed to give correct

bound-state energies that evidence for the Fermi Surface Anomaly emerges. $J_{_{\mbox{$V$}}}$ and the geometry of the optical potentials found for $^{209}_{\mbox{$Bi$}}$ become equal to those needed to explain the high-energy $^{208}_{\mbox{$Pb$}}$ data at about 10.0 MeV. Since $\mbox{dJ}_{_{\mbox{$V$}}}/\mbox{dE}$ for the latter is smaller in magnitude than for $^{209}_{\mbox{$er$}}$ Bi, a change in $\mbox{dJ}_{_{\mbox{$V$}}}/\mbox{dE}$ is clearly indicated near 10.0 MeV. This may effect the extrapolation of higher-energy and charged-particle potentials into the lower-energy neutron domain.

This work supported by the U.S. Department of Energy, Basic Energy Science Programs, under the contract W-31-109-Eng-38.